

MAXIBAN 72 - narasin and nicarbazin granule
Elanco Animal Health Co

*

Net Weight **50 lbs**
(22.68 kg)

Type A Medicated Article

Do not feed undiluted.

Active Drug Ingredients:

Narasin 36 g per lb

Nicarbazin 36 g per lb

For the prevention of coccidiosis in broiler chickens caused by *Eimeria necatrix*, *E. tenella*, *E. acervulina*, *E. brunetti*, *E. mivati*, and *E. maxima*.

Important: Must be thoroughly mixed into feeds before use.

Mixing Directions: (Broiler Chickens): Thoroughly mix the following amounts of Maxiban 72 in one ton of complete feed to provide from 27 to 45 grams per ton each of narasin and nicarbazin. The dosage should be adjusted to meet the severity of the coccidial challenge, which varies with environmental and management conditions. It is recommended that a preblend of the Maxiban 72 in 15-20 lbs of feed be made before incorporating into the total amount of finished feed.

Narasin and Nicarbazin (Grams Per Ton)	Maxiban 72 (Lbs Per Ton of Type C Feed)
54	0.75
72	1.00
90	1.25

Feeding Directions: Feed continuously as the sole ration.

NOT FOR HUMAN USE

WARNING:

Feeds containing Maxiban must be withdrawn 5 days prior to slaughter. When mixing and handling Maxiban 72, use protective clothing, impervious gloves, and a dust mask.

Operators should wash thoroughly with soap and water after handling. If accidental eye contact occurs, immediately rinse thoroughly with water.

Caution: Nicarbazin medicated broilers may show reduced heat tolerance if exposed to high temperature and high humidity. Provide adequate drinking water and ventilation during these periods. Do not allow adult turkeys, horses or other equines access to formulations containing narasin. Ingestion of narasin by these species has been fatal. Do not feed to laying hens.

Store in a cool, dry place.

Not to be used after the date printed on top of bag.

Restricted Drug (California) - Use Only as Directed

NADA # 138-952, Approved by FDA

*Elanco[®], Maxiban[®] and the diagonal color bar are trademarks of Eli Lilly and Company.

Elanco Animal Health

A Division of Eli Lilly and Company

Indianapolis, IN 46285, U.S.A.

To report adverse effects, access medical information, or obtain additional product information, call 1-800-428-4441.

BG5260DEAMB (V02-03-2009)



PRINCIPAL DISPLAY PANEL – 50 LBS BAG

ELANCO* AF1372

For Use In Broiler Chicken Feeds Only

Maxiban® 72

Narasin and Nicarbazine

Net Weight **50 lbs** (22.68 kg)



ELANCO* AF1372

For Use In Broiler
Chicken Feeds Only
Maxiban®
Narasin and
Nicarbazine
72

Net Weight **50 lbs**
(22.68 kg)

Type A Medicated Article

Do not feed undiluted.

Active Drug Ingredients: Narasin 36 g per lb
Nicarbazine 36 g per lb

For the prevention of coccidiosis in broiler chickens caused by *Eimeria necatrix*, *E. tenella*, *E. acervulina*, *E. brunetti*, *E. nivalis*, and *E. maxima*.

Important: Must be thoroughly mixed into feeds before use.

Mixing Directions: (Broiler Chickens): Thoroughly mix the following amounts of Maxiban 72 in one ton of complete feed to provide from 27 to 45 grams per ton each of narasin and nicarbazine. The dosage should be adjusted to meet the severity of the coccidial challenge, which varies with environmental and management conditions. It is recommended that a preblend of the Maxiban 72 in 15-20 lbs of feed be made before incorporating into the total amount of finished feed.

Narasin and Nicarbazine (Grams Per Ton)	Maxiban 72 (Lbs Per Ton of Type C Feed)
54	0.75
72	1.00
90	1.25

Feeding Directions: Feed continuously as the sole ration.

NOT FOR HUMAN USE

Warning: Feeds containing Maxiban must be withdrawn 5 days prior to slaughter. When mixing and handling Maxiban 72, use protective clothing, impervious gloves, and a dust mask. Operators should wash thoroughly with soap and water after handling. If accidental eye contact occurs, immediately rinse thoroughly with water.

Caution: Nicarbazine medicated broilers may show reduced heat tolerance if exposed to high temperature and high humidity. Provide adequate drinking water and ventilation during these periods. Do not allow adult turkeys, horses or other equines access to formulations containing narasin. Ingestion of narasin by these species has been fatal. Do not feed to laying hens.

Store in a cool, dry place.

Not to be used after the date printed on top of bag.

Restricted Drug (California) - Use Only as Directed

NADA # 138-952, Approved by FDA

*Elanco® Maxiban® and the diagonal color bar are trademarks of Eli Lilly and Company.

Elanco Animal Health
A Division of Eli Lilly and Company
Indianapolis, IN 46285, U.S.A.

To report adverse effects, access medical information,
or obtain additional product information, call 1-800-428-4441.

BGS260DEAMB (V02-03-2009)



ELANCO* AF1372

Maxiban®
72

BGS260DEAMB

ELANCO* AF1372

Sólo para uso en alimento
de Pollos de Engorde
Maxiban®
Narasin y
Nicarbazine
72
Peso neto: **50 libras**
(22.68 kg)

Artículo Medicado Tipo A

No administrar sin diluir

Principios Activos: Narasina 36 g per lb (80 g per kg)
Nicarbazine 36 g per lb (80 g per kg)

Para la prevención de coccidiosis en pollos de engorde causada por *Eimeria necatrix*, *E. tenella*, *E. acervulina*, *E. brunetti*, *E. nivalis* y *E. maxima*.
Importante: Debe mezclarse bien con el alimento antes de usarse.
Instrucciones para la mezcla: (Pollos de engorde): Mézclase completamente las siguientes cantidades de Maxiban 72 en una tonelada de alimento completo para proporcionar de 27 a 45 g/tonelada corta (30 a 50 g/ton Métrica), tanto de narasina como de nicarbazine. La dosis debe ajustarse de acuerdo con la gravedad del desafío por coccidias, el cual varía según las condiciones ambientales y de manejo. Se recomienda hacer una premezcla de Maxiban 72 en 15 a 20 libras (7 a 10 kilos) de alimento antes de incorporarlo a la cantidad total de alimento en preparación.

Narasina + Nicarbazine (g/ton)		Maxiban 72 Dosis	
Corta	Métrica	libra corta	g/ton Métrica
54	60	0.75	375
72	80	1.00	500
90	100	1.25	625

Instrucciones de uso: Administre continuamente como ración única.
NO USAR EN SERES HUMANOS

Advertencia: El suministro de alimento conteniendo Maxiban deberá ser retirado 5 días antes del sacrificio. Deberá usarse ropa protectora, guantes impermeables y mascarilla antipolvo al mezclar o manipular el producto. Los operadores deberán lavarse cuidadosamente con agua y jabón después de manejar el producto. Si se produce contacto accidental con los ojos, lavar inmediatamente con agua abundante.

Precaución: Los pollos tratados con nicarbazine pueden presentar menor tolerancia al calor si son expuestos a temperaturas y humedad elevadas. Proveer agua de bebida y ventilación adecuadas durante estos periodos. No permita que pavos adultos, caballos u otros equinos tengan acceso a raciones que contengan narasina ya que la ingestión de narasina por parte de estas especies es fatal. No administrar a gallinas de postura.

Consérvese en un lugar fresco y seco.

No usar después de la fecha impresa en la parte superior del saco.
Medicamento Restringido (California) - Use Solo Como se Indica.
NADA# 138-952, Aprobado por FDA.

*Elanco® Maxiban® y la barra diagonal de color son marcas registradas de Eli Lilly and Company.

Elanco Animal Health • Una División de Eli Lilly and Company
Indianapolis, IN 46285, Estados Unidos de América.

Para reportar efectos adversos, obtener información
medica o información adicional del producto,
llame al 1-800-428-4441.

BGS260DEAMB (V02-03-2009)

